

**10<sup>th</sup> IAPC Meeting**  
**Tenth World Conference on**  
**Physico-Chemical Methods in Drug Discovery**  
**&**  
**Sixth World Conference on ADMET and DMPK**



**Book of**  
**Abstracts**



September 4-6, 2023 :: Belgrade, Serbia

## **10<sup>th</sup> IAPC Meeting**

*Tenth World Conference on Physico-Chemical Methods in Drug Discovery  
&*

*Sixth World Conference on ADMET and DMPK  
Belgrade, Serbia, September 4-6, 2023*

## **Book of Abstracts**

*Organized by*

***International Association of Physical Chemists***

*&*

***Faculty of Chemistry, University of Belgrade, Serbia***

*Published by*

***International Association of Physical Chemists***

*E-mail: [office@iapchem.org](mailto:office@iapchem.org), URL: <http://www.iapchem.org>*

For Publisher

**Zoran Mandić**

Editor

**Tatjana Verbić & Zoran Mandić**

Design, page making and computer layout

**Aleksandar Dekanski**

On Line version only

## The Scientific and Organizing Committee:

**Tatjana Verbić**, Conference Chair  
University of Belgrade, Serbia,

**Alex Avdeef**  
ADME Research, New York, USA

**Kiyohiko Sugano**  
Ritsumeikan University, Osaka, Japan

**Kin Tam**  
University of Macau, Macau

**Zoran Mandić**  
University of Zagreb, Croatia

**Klara Valko**  
Biomimetic chromatography Ltd. UK

**Godefridus J. Peters**  
Amsterdam University Medical Centers, The Netherlands

**Hong Wan**  
WHDeX Consulting AB, Sweden

## Local Organizing Committee

**Tatjana Verbić**, Conference Chair  
University of Belgrade, Faculty of Chemistry, Serbia

**Goran Roglić**  
University of Belgrade, Faculty of Chemistry, Serbia

**Ilija Cvijetić**  
University of Belgrade, Faculty of Chemistry, Serbia

**Miloš Pešić**  
University of Belgrade, Faculty of Chemistry, Serbia

**Olivera Marković**  
University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

**Aleksandar Dekanski**  
University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Serbia

**Marija Popović Nikolić**  
University of Belgrade, Faculty of Pharmacy, Serbia

P 08

## Study of ionization of montelukast in differently charged micellar solutions as biomembrane mimetic systems

Marija Popović-Nikolić<sup>1</sup>, Slavica Oljačić<sup>1</sup>, Katarina Nikolić<sup>1</sup>, Gordana Popović<sup>2</sup>

<sup>1</sup>*Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, Belgrade, Serbia*

<sup>2</sup>*Department of General and Inorganic Chemistry, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, Belgrade, Serbia*

Montelukast is a leukotriene receptor antagonist indicated for asthma prophylaxis in adults as well as in pediatric patients 6 months of age and older. Because it is associated with numerous side effects, including neuropsychiatric events, it is very important to monitor its pharmacologic behavior when administered chronically. To gain better insight into the pharmacological properties of ionizable drugs, their physicochemical properties should be studied under conditions more similar to physiological, such as micellar solutions of surfactants as biomembrane mimetic systems. Montelukast is an ampholyte with one acidic (carboxyl) and one basic (quinoline nitrogen) group. In this study the effects of micellar solutions of differently charged surfactants (anionic SDS, cationic CTAB, and nonionic TX-100) on protolytic equilibria of montelukast were investigated potentiometrically. The solutions ( $5 \times 10^{-4}$  M) were titrated with standard NaOH solution (0.1017 M) at a constant ionic strength (0.1 M NaCl) and a temperature 25°C. Experimental data were analyzed using the Hyperquad program. Due to poor water solubility, the  $pK_a$  values defining the ionization in water ( $pK_{a1}=4.07$ ,  $pK_{a2}=5.49$ ), were obtained indirectly by extrapolation from the  $pK_a^*$  values determined potentiometrically in the different methanol-water mixtures (40%, 50%, and 55% wt/wt). The  $pK_a$  values in 0.01M micellar solutions were determined without the use of cosolvent. Micelles contributed to the shift in protolytic equilibria of montelukast, anionic  $\Delta pK_a$  up to +1.20, cationic  $\Delta pK_a$  up to +0.27, and nonionic  $\Delta pK_a$  up to +0.98. More pronounced effects are observed on the ionization of carboxyl group than quinoline nitrogen. A change in the distribution of equilibrium forms in a relation to pure water, can be expected in physiological conditions, in interactions of montelukast with polar or charged biomolecules.

**Acknowledgments:** This research was funded by the Ministry of Science, Technological Development and Innovation, Republic of Serbia through Grant Agreement with University of Belgrade-Faculty of Pharmacy No: 451-03-47/2023-01/ 200161. The authors thank the COST Actions CA18240 and CA18133 for support.