

BIOCHEMISTRY AND METABOLISM OF VITAMIN D

BIOHEMIJA I METABOLIZAM VITAMINA D

Snežana Jovičić¹, Svetlana Ignjatović², Nada Majkić-Singh²¹Center for Medical Biochemistry, Clinical Center of Serbia, Belgrade, Serbia²Center for Medical Biochemistry, Clinical Center of Serbia and Faculty of Pharmacy, University of Belgrade, Belgrade, Serbia

Summary: Vitamin D is not technically a vitamin, since it is not an essential dietary factor. It is rather a prohormone produced photochemically in the skin from 7-dehydrocholesterol. Vitamin D and its metabolites may be categorized as either cholecalciferols or ergocalciferols. Cholecalciferol (vitamin D₃) is the parent compound of the naturally occurring family and is produced in the skin from 7-dehydrocholesterol on exposure to the ultraviolet B portion of sunlight. Vitamin D₂ (ergocalciferol), the parent compound of the other family, is manufactured by irradiation of ergosterol produced by yeasts and its potency is less than one-third of vitamin D₃'s potency. The steps in the vitamin D endocrine system include the following: 1) the photoconversion of 7-dehydrocholesterol to vitamin D₃ in the skin or dietary intake of vitamin D₃; 2) metabolism of vitamin D₃ by the liver to 25-hydroxyvitamin-D₃ [25(OH)D₃], the major form of vitamin D circulating in the blood compartment; 3) conversion of 25(OH)D₃ by the kidney (functioning as an endocrine gland) to the hormone 1,25-dihydroxyvitamin D₃ [1,25(OH)₂D₃]; 4) systemic transport of the dihydroxylated metabolite 1,25(OH)₂D₃ to distal target organs; and 5) binding of 1,25(OH)₂D₃ to a nuclear receptor (VDR) at target organs, followed by generation of appropriate biological responses. The activation of vitamin D to its hormonal form is mediated by cytochrome P450 enzymes. Six cytochrome P450 (CYP) isoforms have been shown to hydroxylate vitamin D. Four of these, CYP27A1, CYP2R1, CYP3A4 and CYP2J3, are candidates for the enzyme vitamin D 25-hydroxylase that is involved in the first step of activation. The highly regulated, renal enzyme 25-hydroxyvitamin D-1 α -hydroxylase contains the component CYP27B1, which completes the activation pathway to the hormonal form 1,25(OH)₂D₃. A five-step inactivation pathway from

Kratak sadržaj: Vitamin D nije pravi vitamin, odnosno nije esencijalni dijetetski faktor, već je pre prohormon koji nastaje fotohemijskom reakcijom u koži iz 7-dehidroholesterol. Vitamin D i njegovi metaboliti mogu da se kategorizuju kao holekalciferoli ili ergokalciiferoli. Holekalciferol (vitamin D₃) je polazno jedinjenje za familiju koja se nalazi u prirodi i produkuje se u koži iz 7-dehidroholesterol. Vitamin D₂ (ergokalciiferol), polazno jedinjenje druge familije, nastaje radijacijom ergosterola koga produkuju kvasci i ima samo jednu trećinu aktivnosti vitamina D₃. Faze u endokrinom sistemu vitamina D su: 1) fotokonverzija 7-dehidroholesterol u vitamin D₃ u koži ili unos vitamina D₃ hranom; 2) metabolizam vitamina D₃ u jetri do 25-hidroksivitamina D₃ [25(OH)D₃], glavnog oblika vitamina D u cirkulaciji; 3) konverzija 25(OH)D₃ u bubregu (koji ovde funkcioniše kao endokrina žlezda) do hormona 1,25-dihidroksivitamin D₃ [1,25(OH)₂D₃]; 4) sistemski transport dihidroksi-metabolita do distalnih ciljnih organa; i 5) vezivanje 1,25(OH)₂D₃ za nuklearni receptor (VDR) u ciljnim organima, što prati odgovarajući biološki odgovor. Aktivacija vitamina D do hormonskog oblika je posredovana citohrom P450 enzimima. Pokazano je da šest izoformi citohroma P450 (CYP) učestvuje u hidr oksilaciji vitamina D. Za četiri od njih, CYP27A1, CYP2R1, CYP3A4 i CYP2J3, se pretpostavlja da imaju aktivnost 25-hidroksilaze koja učestvuje u prvom koraku aktivacije. Renalni enzim, 25-hidroksivitamin D-1 α -hidroksilaza sastrugoregulisanom aktivnošću, predstavlja CYP27B1, koji završava aktivaciju do hormonskog oblika 1,25(OH)₂D₃. Proces inaktivacije, koji se sastoji iz pet stupnjeva od 1,25(OH)₂D₃ do kalcitrične kiseline, obavlja jedan multifunkcionalni CYP, CYP24A1, čija je transkripcija indukovana u ciljnim ćelijama.

Address for correspondence:

Snežana Jovičić
Center for Medical Biochemistry
Clinical Center of Serbia
Višegradska 26, 11000 Belgrade
Tel/Fax: + 381 11 361 56 31
e-mail: hionati@gmail.com

1,25(OH)₂D₃ to calcitric acid is attributed to a single multifunctional CYP, CYP24A1, which is transcriptionally induced in vitamin D target cells by the action of 1,25(OH)₂D₃. An additional key component in the operation of the vitamin D endocrine system is the plasma vitamin D binding protein (DBP), which carries vitamin D₃ and its metabolites to their metabolism and target organs. DBP is a specific, high-affinity transport protein. It is synthesized by the liver and circulates in great excess, with fewer than 5% of the binding sites normally occupied. 1,25(OH)₂D₃ acts as a ligand for a nuclear transcription factor, vitamin D receptor – VDR, which like all other nuclear receptors, regulates gene transcription and cell function. The widespread presence of VDR, and the key activating (1 α -hydroxylase, CYP27B1) and inactivating (24-hydroxylase, CYP24A1) enzymes in most mammalian cells means that the cells in these tissues have the potential to produce biological responses, depending on the availability of appropriate amounts of vitamin D₃. Thanks to this widespread presence of elements of vitamin D endocrine system, its biological features are being recognized outside bone tissue, i.e. calcium and phosphate metabolism.

Keywords: vitamin D, vitamin D receptor, vitamin D binding protein, cytochrome P450

Introduction

Vitamin D₃ is not a genuine vitamin, in the sense that it is not an essential element of diet, but it is more of a steroid prohormone produced through photochemical reaction in the skin from 7-dehydrocholesterol. The metabolic action is carried out through its hormonal form, 1 α ,25-dihydroxyvitamin D₃ [1,25(OH)₂D₃], after binding to a nuclear receptor (vitamin D receptor, VDR) which regulates the transcription of a number of target genes in a variety of vitamin D target cells. The presence of activating and inactivating enzymes of vitamin D, as well as VDR in almost all mammalian cells, together with the observation that approximately 3% of the mouse and human genome is regulated via the vitamin D pathway, indicates that it is essential for life in higher animals (1). Besides the well established central role in calcium homeostasis, this pluripotent hormone has additional biological actions in the adaptive and innate immune system, insulin secretion by the pancreatic β cells, multifactorial heart functioning and blood pressure regulation, brain and fetal development (2). In accordance, the all growing epidemiological data support the role that vitamin D might play in controlling the risk of many chronic illnesses, including common cancers, myopathy, autoimmune disease, diabetes and the metabolic syndrome, infections, and cardiovascular disease (1, 3).

In this article we summarize the current knowledge of the metabolism and biochemistry of vitamin D, including biological activation and inactivation, mechanisms of action and regulation.

ma dejstva vitamina D posredstvom 1,25(OH)₂D₃. Dodatna ključna komponenta u dejstvu vitamin D endokrinog sistema je vitamin D vezujući protein u plazmi (DBP), koji transportuje vitamin D₃ i njegove metabolite do ciljnih i organa gde se odvija njihov metabolizam. DBP je specifičan transportni protein velikog afiniteta. Sintetiše se u jetri i cirkuliše u velikom višku, sa zasićenjem vezujućih mesta manjim od 5%. 1,25(OH)₂D₃ deluje kao ligand nuklearnog transkripcionog faktora, VDR, koji reguliše transkripciju gena i funkciju ćelija. široka rasprostranjenost VDR i ključnih enzima aktivacije (1 α -hidroksilaza, CYP27B1) i inaktivacije (24-hidroksilaza, CYP24A1) u većini ćelija sisara znači da ćelije u ovim tkivima imaju potencijal za proučavanje bioloških odgovora, zavisno od raspoloživosti dovoljnih količina vitamina D₃. Zahvaljujući rasprostranjenosti elemenata endokrinog sistema vitamina D, njegove biološke osobine se prepoznaju i izvan koštanog sistema, odnosno metabolizma kalcijuma i fosfora.

Ključne reči: vitamin D, receptor za vitamin D, vitamin-D vezujući protein, citohrom P450

Structure and synthesis of vitamin D

Vitamin D is a secosteroid with broken 9,10 carbon-carbon bond in B ring of the cyclopentanoperhydrophenanthrene structure. Vitamin D and its metabolites may be classified as either cholecalciferols or ergocalciferols. Cholecalciferol (vitamin D₃) is the parent compound of the naturally occurring family and is produced in the skin from 7-dehydrocholesterol on exposure to the ultraviolet B portion of sunlight. Beside the photosynthesis in the skin, vitamin D₃ can also be introduced with certain types of foods, including fatty fish, fish liver oils, and egg yolk. Vitamin D₂ (ergocalciferol) is manufactured by irradiation of ergosterol produced by yeasts. Vitamin D₂ differs from vitamin D₃ by the double bond between carbon 22 and carbon 23 and a methyl group on carbon 24. When vitamin D or its metabolites are written without a subscript, both families are included (4). Vitamin D₂ has only one third of vitamin D₃'s potency regarding its biological actions (5). Structure of vitamins D₃ and D₂ is represented in Figure 1.

Metabolism

Vitamin D₃ has no known biological function in its native form. It must be metabolized first to 25-hydroxyvitamin D₃ [25(OH)D₃] in the liver and then to 1,25(OH)₂D₃ by the kidney. In studies of vitamin D metabolism some 37 vitamin D₃ metabolites have been isolated and chemically characterized (6). All of these metabolites can be systematized into three basic pathways of vitamin D₃ metabolism: (a) the main two-step activation pathway in liver and kidney that produces 1,25(OH)₂D₃; (b) an inducible carbon-




Figure 1 Structure of vitamin D₃ (cholecalciferol) and vitamin D₂ (ergocalciferol) and their precursors (4).

24 oxidation pathway in vitamin D target cells for inactivating 25(OH)D₃ to 24,25-dihydroxyvitamin D₃ [24,25(OH)₂D₃] and 1,25(OH)₂D₃ to calcitroic acid; (c) and not completely elucidated 26,23-lactone pathway for converting both 25(OH)D₃ and 1,25(OH)₂D₃ to lactone products (7). Vitamin D₂ undergoes 25- and 1 α -hydroxylation steps by the same enzymes, which produces an analogous form of 1,25(OH)₂D₂, after the conversion into 25(OH)D₂ (8).

The sequencing of the human genome has led to understanding that among a pool of 60 total cytochrome P450s (CYPs) there are three known and possibly other vitamin D-related CYPs linked to the metabolism of vitamin D. The key enzymes in vitamin D metabolism are the hepatic vitamin D-25-hydroxylase (CYP27A1 and CYP2R1), renal 25-hydroxyvitamin D-1 α -hydroxylase (CYP27B1), and 25-hydroxyvitamin D-24-hydroxylase (CYP24A1) (9).

Vitamin D-25-hydroxylase

Several CYPs are able to catalyze the 25-hydroxylation of vitamin D₃ *in vitro*. No meaningful regula-

tion of CYP27A1 by vitamin D or calcium homeostatic hormones has been reported in the literature, why it is probably not the physiologically relevant 25-hydroxylase. CYP27A1 does not 25-hydroxylate vitamin D₂ and the efficient 25-hydroxylation of vitamin D₂ has been reported in most mammals *in vivo*. This enzyme has an essential role in cholesterol and bile acid metabolism and its deficiency does not affect serum levels of vitamin D metabolites. Genetic mutations of CYP27A1 in humans cause cerebrotendinous xanthomatosis, a disorder of bile acid and lipid metabolism that sometimes presents with low 25(OH)D levels and a type of osteoporosis (7).

On the other hand, microsomal CYP2R1 is thought to play a major role due to the absence of sex and species differences and catalytic activity toward both vitamin D₂ and D₃. CYP2R1 is regiospecific to the C-25 position of a secosteroid in contrast to other polyfunctional CYP enzymes with vitamin D 25-hydroxylase activity (CYP27A1, CYP2C11, and CYP3A4). Moreover, mutation in the human *Cyp2r1* gene results in vitamin D-dependent rickets, the only type of rickets that is caused by 25-hydroxylase deficiency. Furthermore, CYP2R1 expression in human

tissues is ubiquitous, which supports the findings that the ability to form $25(\text{OH})\text{D}_3$ is almost unchanged in patients with liver disorders (10).

25-hydroxyvitamin D-1 α -hydroxylase

Mitochondrial 25-hydroxyvitamin D-1 α -hydroxylase (CYP27B1) has a central role in calcium homeostasis. The main control of calcium homeostasis is the calcium-sensing receptor in the parathyroid cell, which regulates the secretion of parathyroid hormone (PTH). In turn, PTH is principal activator of 25-hydroxyvitamin D-1 α -hydroxylase gene expression, which represents the mechanism needed to control plasma concentration of $1,25(\text{OH})_2\text{D}$ (11). This regulation is independently amplified by low calcium and phosphorus signals (12). By maintaining tighter regulation of the concentration of $1,25(\text{OH})_2\text{D}$ and thereby giving rise to appropriate transcriptional activation of the genes involved in calcium and phosphorus transport and cell differentiation, the 25-hydroxyvitamin D-1 α -hydroxylase plays a vital role in vitamin D signaling (7).

It has been shown, however, that CYP27B1 is expressed in various extrarenal sites around the body including the keratinocyte, lung, colon, and macrophages (1). This indicates that extrarenal 25-hydroxyvitamin D-1 α -hydroxylase has an autocrine or paracrine role in specific tissue differentiation (13). This was demonstrated by the work of Dusso et al. (14) who have shown that cytokines such as interferon- γ , not PTH, upregulate expression of CYP27B1 in the macrophage. The concept of extrarenal 25-hydroxyvitamin D-1 α -hydroxylase has not only physiological implications but also pathological ones. For example, sarcoidosis is a granulomatous condition that often involves hypercalciuria and eventually hypercalcemia caused by the overproduction of $1,25(\text{OH})_2\text{D}$ in sarcoid macrophages (7).

25-hydroxyvitamin D-24-hydroxylase

Mitochondrial 25-hydroxyvitamin D-24-hydroxylase (CYP24A1) is also regulated by $1,25(\text{OH})_2\text{D}$, but in opposite direction to 25-hydroxyvitamin D-1 α -hydroxylase (CYP27B1) owing to the transcriptional upregulation in the CYP24A1 promoter. The outcome is induction of CYP24A1 in all vitamin D target cells, which provides exquisite attenuation of the hormonal signal in the individual target cell when the gene transcriptional upregulation of $1,25(\text{OH})_2\text{D}$ need to be turned off (7). *Cyp24a1* knockout mice confirm the catabolic role for CYP24A1, because they show poor viability with 50% lethality, showing hypercalcemia, with marked difficulty in excreting a bolus of [1β - ^3H] $1\alpha,25$ - $(\text{OH})_2\text{D}_3$ and isolated cultured keratinocytes from these mice fail to synthesize calcitroic acid, as opposite to heterozygous and wild-

type mice (15). In humans, mutations in *Cyp24a1* with a complete loss of function, cause idiopathic infantile hypercalcemia (16).

C-24 oxidation represents the predominant catabolic pathway for $1,25(\text{OH})_2\text{D}_3$ in which sequential C-24 hydroxylation, C-24 ketonization, and C-23 hydroxylation, followed by oxidative cleavage to form tetranor- $1\alpha, 23$ -dihydroxyvitamin D_3 and calcitroic acid, liberate an inactive, water-soluble product that is excreted in bile (9). The secondary pathway involves C-26 hydroxylation, C-23 hydroxylation, and lactonization to form (23S,25R)- $1,25(\text{OH})_2\text{D}_3$ -26,23-lactone. Both pathways have also been shown to participate in the metabolism of $25(\text{OH})\text{D}_3$ (17). Studies of the recombinant CYP24A1 protein have revealed that CYP24A1 uses multicatalytic activity, facilitating sequential oxidation of C-23, C-24 and C-26 hydroxylation and side-chain cleavage (9).

Mechanisms of action

Mechanism of action of vitamin D_3 , through its hormonal form, $1,25(\text{OH})_2\text{D}_3$ involves nuclear receptor (VDR) that regulates the transcription of several target genes in a variety of vitamin D target cells that are primarily involved in calcium homeostasis and cell differentiation (18). These regulatory functions are performed by several specific proteins included in vitamin D signal transduction system, which constitutes of VDR and its associated transcriptional coactivators, plasma transport protein (vitamin D binding protein, DBP), the activating CYPs (CYP2R1, CYP27A1, and CYP27B1), and catabolic CYP24A1 (described in previous section) (19).

Vitamin D binding protein

Because of its lipophilic nature vitamin D_3 requires a protein carrier for solubility in plasma. Its mono-, di-, and tri-hydroxylated metabolites show progressively increasing polarity, culminating in the water soluble biliary form of calcitroic acid. After the absorption from the gut, vitamin D enters the circulation on chylomicrons first, and it is only slowly transferred to DBP, for which it has low affinity, between 10^{-5} and 10^{-7} mol/L. The difference between the transport of dietary and vitamin D synthesized in the skin is that the latter is mainly bound to DBP (20). The consequence of chylomicron transport of dietary vitamin D is the possibility of uptake by peripheral tissues, such as adipose tissue and muscle, due to the action of lipoprotein lipase. The liver takes up remaining vitamin D from the chylomicron remnant and quickly removes it from the bloodstream. The loss into tissue and liver pools explains the short plasma half-life, ≈ 4 – 6 h, and the whole-body half-life ≈ 2 months of vitamin D (19).

In the liver vitamin D converts into 25(OH)D due to the action of microsomal CYP2R1 or mitochondrial CYP27A1, neither of which is subject to tight regulation (21). 25(OH)D quickly enters the plasma pool that constitutes the predominant pool of vitamin D in the body, with a capacity of $\approx 4.5 \mu\text{mol/L}$. 25(OH)D₃ and 25(OH)D₂ have a strong affinity for DBP at $5 \times 10^{-8} \text{ mol/L}$ why 25(OH)D₃ has a half-life of 15 days in the circulation. The normal circulating level of 25(OH)D in the blood is only 25–200 nmol/L, indicating that ligand only occupies 2–5% of DBP in the physiologic state (19).

The dihydroxy- metabolites have different affinities for DBP: 25(OH)D₃-26,23-lactone binding 3–5 times as tightly as 25(OH)D₃ and inactive metabolites, 24,25(OH)₂D₃ and 25,26(OH)₂D₃, bind with equal affinity as 25(OH)D₃. The active form 1,25(OH)₂D₃ has a half-life of 10–20 h depending on the state of the highly inducible catabolic machinery. The accumulation of these metabolites in the bloodstream is mainly a function of their affinities for DBP, but the rate of synthesis and degradation also plays a partial role (19).

Vitamin D receptor

The steroid hormone 1,25(OH)₂D₃, like other steroid hormones, generates biological responses by regulating gene transcription (genomic responses) and by activating a variety of signal transduction pathways at or near plasma membrane (nongenomic rapid responses) (22). The genomic responses are due to stereospecific interaction of 1,25(OH)₂D₃ with its nuclear receptor, VDR. The primary amino acid sequence of the VDR consists, like in other steroid hormone receptors, of 6 functional domains: the variable regions, DNA binding, the hinge region, the ligand binding region, and the transcriptional activation domain (2). Formation of the ligand-receptor complex results in conformational changes in the receptor protein, which allow the ligand-receptor complex to specifically interact with many proteins from the transcriptional machinery. It is estimated that the VDR can regulate the expression of as many as 500 of the 20 488 genes in the human genome (23). The large number of VDR-regulated genes undoubtedly reflects the consequence of the distribution of both the VDR and 25(OH)D₃-1 α -hydroxylase to many organs.

»Rapid« or nongenomic responses provoked by 1,25(OH)₂D₃ are mediated through the interaction of the hormone with a receptor located on the cell's external membrane. This membrane receptor is the classic VDR, found in nucleus and cytosol and associated with caveolae in the plasma membrane of a variety of cells (24). Caveolae are flask-shaped membrane invaginations enriched in sphingolipids and cholesterol that are commonly found in a wide variety of cells (25). Using VDR knockout and wild-type mice, it was found that rapid modulation of osteoblast

ion channel responses by 1,25(OH)₂D₃ require the presence of a functional vitamin D nuclear and caveolae receptor (26).

Although VDR is expressed in many cells around the body, differences in tissue-, differentiation stage- and gene-specific transcription factors present at the vitamin D-dependent genes allow wide variability in the range of genes that are modulated in each tissue at any given time. Even the direction of the effect of 1,25(OH)₂D₃ on gene transcription, in the sense whether it causes upregulation or downregulation, is gene-specific. For example, various Ca-homeostatic genes (e.g. calbindins, Ca-channel proteins, osteocalcin, osteopontin, RANKL genes) are upregulated, whereas others (e.g. collagen and pro-pro-PTH genes) are downregulated by 1,25(OH)₂D₃ (7).

VDR is 1,25(OH)₂D₃-dependent transcription factor that controls gene expression by heterodimerizing with retinoid X receptors (RXRs) and associating specifically with VDR responsive elements (VDRE) in target genes. Sequence and promoter analysis of several 1,25(OH)₂D₃-regulated genes have led to the identification of VDREs, short DNA sequences to which the VDR-RXR heterodimer binds and subsequently exerts its influence on transcription. Some VDREs have been identified in genes that are known to be transcriptionally activated by the 1,25(OH)₂D₃ hormone including osteocalcin and osteopontin (expressed in bone osteoblasts), β_3 integrin (found in bone osteoclasts and macrophages), calbindin-D_{28k} (from kidney) and p21 (an inhibitor of cyclin dependent kinase, Cdk, in many tissues) (12).

The activation of the ligand-bound VDR in the intestine, bone, kidney, and parathyroid gland cells results in the maintenance of normal serum calcium and phosphorus levels and their related effects on mineralization and turnover of bone (27). However, 1 α -hydroxylase is also present in cells of several extrarenal tissues such as skin, bone, prostate, and many immune cells. The enzyme here is identical to the one expressed in the kidney, but its expression is regulated by immune signals instead of mediators of bone and mineral homeostasis (28). Therefore, the potential actions of 1,25(OH)₂D₃ via its nuclear VDR extend far beyond the bone mineral homeostasis. High local 1,25(OH)₂D₃ concentrations may, independently from serum concentrations, manifest an autocrine and paracrine function since the VDR is widely present in many different cells and tissues (29). For example, one major target for 1,25(OH)₂D₃ is the immune system, where mediated by VDR it suppresses IL-1 to IL-6 and interferon- γ *in vitro*. Moreover, documented *in vivo* immunomodulatory actions of the hormone are reduced macrophage and lymphocyte function in vitamin D-deficient rats. 1,25(OH)₂D₃ functions as a general suppressor of the immune system, especially of Th1 cells, suggesting its potential usefulness in the treatment of autoimmune disorders. In central nervous system, besides

immunosuppression, $1,25(\text{OH})_2\text{D}_3$ has been shown to induce expression of a number of neurotrophic hormones from degenerative processes. Together with the expression of calbindin- $\text{D}_{28\text{k}}$ which exhibits antiapoptotic effect, $1,25(\text{OH})_2\text{D}_3$ may have a possible role in therapeutic intervention for neurodegenerative disorders. Similarly, $1,25(\text{OH})_2\text{D}_3$ affects the maturation and function of certain normal and neoplastic cells (e.g. mammary, prostate, and colon), which may be related to the ability of liganded VDR to arrest cells at the G1 stage by influencing cell cycle regulatory proteins, such as p21 and p27, to control cell growth transcription factors, such as *c-myc* and *c-fos*, or to elicit apoptosis by downregulating *Bcl-2*. $1,25(\text{OH})_2\text{D}_3$ also reportedly affects several major endocrine processes, such as TRH/TSH action and pancreatic insulin secretion (12).

Conclusion

In the era of worldwide vitamin D deficiency, the new roles of vitamin D, beyond well established bone

and mineral metabolism, are being recognized. Enzymes involved in the activation of components of this endocrine system, together with elements of catabolic reactions, as well as nuclear receptors, and aspects of autocrine and paracrine actions, have essential roles not only in preserving bone and mineral homeostasis, but in regulation of numerous processes of specific cell differentiation and proliferation. The knowledge of these mechanisms of action of vitamin D endocrine system can be used to help the diagnosis and treatment of diseases involving specific organs and tissues which respond to actions of vitamin D.

Acknowledgments. This study was conducted as a part of the Project No. ON175036, financially supported by the Ministry of Education and Science of Republic of Serbia.

Conflict of interest statement

The authors stated that there are no conflicts of interest regarding the publication of this article.

References

- Bouillon R, Bischoff-Ferrari H, Willet W. Vitamin D and health: perspectives from mice to man. *J Bone Miner Res* 2008; 23(7): 974–9.
- Norman AW. From vitamin D to hormone D: fundamentals of the vitamin D endocrine system essential for good health. *Am J Clin Nutr* 2008; 88(suppl): 491S–9S.
- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007; 357(3): 266–81.
- Endres DB, Rude RK. Mineral and bone metabolism. In: Burtis CA, Ashwood ER, Bruns DE, editors. *Tietz Textbook of Clinical Chemistry*. 4th ed. Philadelphia: WB Saunders, 2006: 1929–6.
- Armas LAG, Hollis BW, Heaney RP. Vitamin D_2 is much less effective than vitamin D_3 in humans. *J Clin Endocrinol Metab* 2004; 89:5387–91.
- Bouillon R, Okamura WH, Norman AW. Structure-function relationships in the vitamin D endocrine system. *Endocr Rev* 1995; 16: 200–57.
- Prosser DE, Jones G. Enzymes involved in the activation and inactivation of vitamin D. *Trends in biochemical sciences* 2004; 29(12): 664–73.
- Haddad JG, Matsuoka LY, Hollis BW, Hu YZ, Wortsman J. Human plasma transport of vitamin D after its endogenous synthesis. *J Clin Invest* 1993; 91: 2552–5.
- Masuda S, Byford V, Arabian A, Sakai Y, Demay MB, St-Arnaud E, et al. Altered pharmacokinetics of $1,25$ -dihydroxyvitamin D_3 and 25 -hydroxyvitamin D_3 in the blood and tissues of the 25 -hydroxyvitamin D - 24 -hydroxylase (*Cyp24a1*) null mouse. *Endocrinology* 2005; 146(2): 825–34.
- Strushkevich N, Usanov SA, Plotnikov AN, Jones G, Park HW. Structural analysis of CYP2R1 in complex with vitamin D_3 . *J Mol Biol* 2008; 380: 96–106.
- Armbrecht HJ, Boltz MA, Hodam TL. PTH increases renal $25(\text{OH})\text{D}_3$ - 1α -hydroxylase (CYP1 α) mRNA but not renal $1,25(\text{OH})_2\text{D}_3$ production in adult rats. *Am J Physiol Renal Physiol* 2003; 284: F1032–6.
- Haussler MR, Whitfield GK, Haussler CA, Hsieh JC, Thompson PD, Selznick SH, et al. The nuclear vitamin D receptor: biological and molecular regulatory properties revealed. *J Bone Miner Res* 1998; 13: 325–49.
- Bikle DD, Chang S, Crumrine D, Elalieh H, Man MQ, Choi EH, et al. 25 -Hydroxyvitamin D - 1α -hydroxylase is required for optimal epidermal differentiation and permeability barrier homeostasis. *J Invest Dermatol* 2004; 122: 984–92.
- Dusso AS, Kamimura S, Gallieni M, Zhong M, Negrè L, Shapiro S, et al. γ -Interferon-induced resistance to $1,25$ - $(\text{OH})_2\text{D}_3$ in human monocytes and macrophages: a mechanism for the hypercalcemia of various granulomatoses. *J Clin Endocrinol Metab* 1997; 82: 2222–32.
- St-Arnaud R. Targeted inactivation of vitamin D hydroxylases in mice. *Bone* 1999; 25: 127–9.
- Schlingmann KP, Kaufmann M, Weber S, Irwin A, Goos C, John U, et al. Mutations in *CYP24A1* and idiopathic infantile hypercalcemia. *N Engl J Med* 2011; 365(5): 410–21.
- Siu-Cadera ML, Zou L, Ehrlich MG, Schwartz ER, Ishizuka S, Reddy GS. Human osteoblasts in culture metabolize both $1\alpha,25$ -dihydroxyvitamin D_3 and its pre-

- cursor 25-hydroxyvitamin D₃ into their respective lactones. *Endocrinology* 1995; 136: 4195–203.
18. Jones G, Strugnell S, DeLuca HF. Current understanding of the molecular actions of vitamin D. *Physiol Rev* 1998; 78: 1193–231.
 19. Jones G. Pharmacokinetics of vitamin D toxicity. *Am J Clin Nutr* 2008; 88(suppl): 582S–6S.
 20. Haddad JG, Matsuoka LY, Hollis BW, Hu YZ, Wortsman J. Human plasma transport of vitamin D after its endogenous synthesis. *J Clin Invest* 1993; 91: 2552–5.
 21. Jones G, Byford V, West S, et al. Hepatic activation and inactivation of clinically-relevant vitamin D analogs and prodrugs. *Anticancer Res* 2006; 26: 2589–96.
 22. Norman AW, Mizwicki MT, Norman DPG. Steroid hormone rapid actions, membrane receptors and a conformational ensemble model. *Nat Rev Drug Discov* 2004; 3: 27–41.
 23. Carlberg C. Current understanding of the function of the nuclear vitamin D receptor in response to its natural and synthetic ligands. *Recent Results Cancer Res* 2003; 164: 29–42.
 24. Huhtakangas JA, Olivera CJ, Bishio JE, Zanello LP, Norman AW. The vitamin D receptor is present in caveolae-enriched plasma membranes and binds 1,25(OH)₂-vitamin D₃ in vivo and in vitro. *Mol Endocrinol* 2004; 18: 2660–71.
 25. Razani B, Woodman SE, Lisanti MP. Caveolae: from cell biology to animal physiology. *Pharmacol Rev* 2002; 54: 431–67.
 26. Bula CM, Huhtakangas J, Olivera CJ, Bishop JE, Norman AW, Henry HL. Presence of a truncated form of the vitamin D receptor (VDR) in a strain of VDR-knockout mice. *Endocrinology* 2005; 146: 5581–6.
 27. De Luca HF. Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr* 2004; 80: 1689S–96S.
 28. Van Etten E, Mathieu C. Immunoregulation by 1,25-hydroxyvitamin D₃: basic concepts. *J Steroid Biochem Mol Biol* 2005; 97: 93–101.
 29. Janssens W, Lehouck A, Carremans C, Bouillon R, Mathieu S, Decramer M. Vitamin D beyond bones in chronic obstructive pulmonary disease. *Time to act. A J Respir Crit Care Med* 2009; 179: 630–6.

Received: May 21, 2012

Accepted: June 21, 2012