



Children's health risk assessment based on the content of toxic metals Pb, Cd, Cu and Zn in urban soil samples of Podgorica, Montenegro

Procena rizika po zdravlje dece na osnovu sadržaja toksičnih metala Pb, Cd, Cu i Zn u gradskom zemljištu na teritoriji Podgorice, Crna Gora

Boban Mugoša*, Dijana Djurović*, Aleksandra Pirnat*, Zorica Bulat†, Snežana Barjaktarović-Labović‡

*Institute of Public Health, Podgorica, Montenegro; †Faculty of Pharmacy, University of Belgrade, Belgrade, Serbia; ‡Health Center Bar, Bar, Montenegro

Abstract

Background/Aim. Due to their low tolerance to pollutants and hand-to-mouth pathways the health risk is very high in children's population. The aim of this study was to evaluate risk to children's health based on the content of heavy metals in urban soil samples from Podgorica, Montenegro. This study included the investigation of several toxic metals such as Pb, Cd, Cu and Zn in soil samples from public parks and playgrounds. **Methods.** Sampling was conducted in a period October–November, 2012. Based on cluster analysis, soil samples were divided into two groups related to similarity of metal content at examined locations: the group I – near by recreational or residential areas of the city, and the group II – near traffic roads. Concentration of toxic metals, in urban soil samples were determined by a graphite furnace atomic absorption spectrometry (Pb and Cd) and by inductively coupled plasma optical emission spectrometry technique after microwave digestion. Due to exposure to urban soil, non-carcinogenic index hazardous index (HI) for children was estimated using 95th percentile values of total metal concentration. The value of the total

(ingestion, dermal and inhalation) HI is calculated for maximum, minimum and the average concentration of metals for children. **Results.** Mean concentrations of Pb, Cd, Cu and Zn in the surface layer of the studied urban soils were 85.91 mg/kg, 2.8 mg/kg and 52.9 mg/kg and 112.5 mg/kg, respectively. Samples from group II showed higher metal content compared to group I. Urbanization and traffic are the main sources of pollution of the urban soils of Podgorica. Most of the samples (93.5%) had a high Pb content, 12.9% of the samples had a higher content of Cd, while Cu and Zn were within the limits prescribed by national legislation. At one location the level of security for lead is $HI = 0.8$ and very closed to maximum acceptable value of 1. It is probably the result of intensive traffic near by. **Conclusion.** All metals investigated showed relatively higher concentrations at sites that were close to industrial places and high ways. The mean concentrations of Pb and Zn and maximum concentrations of Pb, Cd, and Zn were higher than presented values in the National Regulation.

Key words: metals, heavy; soil; risk assessment; health; child.

Apstrakt

Uvod/Cilj. Zbog niske tolerancije na zagađivače i puta prenošenja ruke-usta, rizik od narušenja zdravlja je veoma visok u dečjoj populaciji. Cilj ovog rada bio je da se izvrši procena rizika po zdravlje dece na osnovu sadržaja teških metala u uzorcima gradskog zemljišta u Podgorici, Crna Gora. Ovo istraživanje je obuhvatilo određivanje koncentracija nekoliko toksičnih metala kao što su Pb, Cd, Cu i Zn u uzorcima zemljišta javnih parkova i dečijih igrališta. **Metode.** Uzorkovanje je sprovedeno tokom oktobra i novembra 2012. godine. Klaster analizom uzorci zemljišta podeljeni su u dve grupe na osnovu sličnosti sadržaja metala na ispitivanim lokacijama: grupa I – uzorci sa mesta u rekreativnim ili stambenim delovima grada i grupa II – uzorci iz parkova i igralište blizu saobraćajnice. Koncentracija metala u uzorci-

ma urbanog zemljišta određena je primenom atomske apsorpcione spektrometrije pomoću grafitne kivete (Pb i Cd) i tehnike optičke emisije spektroskopije sa induktivno kuplovanom plazmom (Cu i Zn) nakon mikrotalasne digestije. Na osnovu stepena izloženosti uticaju gradskog zemljišta, nekancerogeni indeks opasnosti (*hazard index* – HI) za decu izračunat je na 95. procentu vrednosti ukupne koncentracije metala. Vrednost za ukupni (ingestioni, dermalni i inhalacioni) HI izračunata je za maksimalne, minimalne i srednje koncentracije ispitivanih metala za decu. **Rezultati.** Srednja koncentracija toksičnih metala u uzorcima površinskog sloja zemljišta iznosila je 85,91 mg Pb/kg tla, 2,8 mg Cd/kg tla, 52,9 mg Cu/kg tla i 112,5 mg Zn/kg tla. Sadržaj metala bio je značajno veći u uzorcima zemljišta iz grupe II nego u uzorcima grupe I. Ovo jasno ukazuje na izražen i uočljiv uticaj urbanizacije, a naročito saobraćaja, na zagađenje zem-

ljišta. Većina uzoraka (93,5%) imala je povećan sadržaj Pb, 12,9% uzoraka imalo je povećan sadržaj Cd, dok je sadržaj Cu i Zn bio u granicama propisanim nacionalnom regulativom. Na jednoj lokaciji vrednost za nivo bezbednosti za olovo bio je HI = 0,8, vrlo blizu maksimalne prihvatljive vrednosti koja iznosi 1, što je verovatno posledica intenzivnog saobraćaja u neposrednoj blizini ispitivane lokacije.

Introduction

Continual urbanization and industrialization induces metals emissions into the terrestrial environment which may greatly influence human health¹. Samples of soils become a very good diagnostic tool of environmental conditions that influence human health^{2,3}. Chemical composition of soil has been conducted in many studies during the last ten years. Special attention has been devoted to studies on urban park playgrounds. Dermal contact, ingestion and inhalation are the main route of exposure to toxic metals in urban environment^{2,4}. A high concentration of toxic metals in urban soils is an important source of human metal intake. Possibility of exposure to adverse effects of soil ingestion is higher in children than adults². Urban children mainly come in contact with soil in parks and playgrounds. A significant amount of toxic metals children could ingest from soil, dust and air⁵. Due to their low tolerance to pollutants and hand-to-mouth pathways the health risk is very high in this population^{6,7}. So, the control of potentially harmful substances in soil is of high importance and has to be kept at low level in the areas frequented by children⁸.

As heavy metals are nondegradable and there is no known homeostasis mechanism for them, any high level of this pollutant may affect the human health affecting the normal functioning of organs, liver, kidney, central nervous system, bones, etc, or acting as cofactors in other diseases^{9,10}.

The aim of this study was to evaluate risk assessment to children's health based on the content of toxic metals in urban soil samples of Podgorica, Montenegro. This study included the investigation of several toxic metals such as Pb, Cd, Cu and Zn in surface soil samples from public parks, playgrounds and kindergartens of Podgorica. Children health risk due to children's toxic metal exposure from urban soil according to hazardous indices (His) was estimated.

Method

Sampling and analysis

This study presents concentrations of four toxic metals, Pb, Cd, Cu and Zn, in surface soil samples from the city's playgrounds in public parks, playgrounds and kindergartens of Podgorica, the capital of Montenegro. A total of 31 parks and playgrounds from the different location of the city were studied. Sampling was conducted during October and November, 2012. Samples of approximately 500 g weight, from top 10 cm layer, within 20 × 20 cm of surface soil, consisting of three sub-samples, were taken and mixed to obtain a bulk

Zaključak. Ispitivani metali imali su više koncentracije na mestima u blizini industrijske zone i autoputa. Prosečne koncentracije Pb i Zn, kao i maksimalne koncentracije Pb, Cd i Zn bile su više od vrednosti propisanih nacionalnom regulativom.

Ključne reči: metali, teški; zemljište; rizik, procena; zdravlje; deca.

composite sample at each playground. Sampling was conducted near by playground equipment such as swings, slides, etc. Stainless trowel was used for sampling and samples are transferred to the laboratory in plastic bags. Stones and foreign objects were hand-removed, and the samples were air-dried for several days. After drying at room temperature samples were gently crushed and sieved to 2 mm and 1.0 ± 0.01 g was weighed for analysis. Microwave acid digestion based on US EPA 3052 method was used for sample preparation. The concentrations of Pb and Cd were determined by a graphite furnace atomic absorption spectrometry (GFAAS) (240Z AA Agilent Technologies-Netherlands) and Cu and Zn by inductively coupled plasma-optical emission spectrometry (ICP-OES) (AMETEC-SpectroArcos, Germany).

Reagents and standards

All chemicals used through the study were analytical grade chemicals. There was no further purification for preparation of all reagents and calibration standards. Deionized ultra pure water was used with conductivity < 1 μS/cm. Certified metal stock solution of 1,000 mg/L (J.T. Baker) by successive dilution with deionized water was used for preparing standards for calibration. Each sample was carried out in triplicate.

Data analysis and risk assessment

Statistical package (SPSS 17.0 for Windows) was used for statistical data analysis. This software uses the upper limit of the 95 percent confidence interval (95 percent upper confidence limit – UCL) for the mean concentrations for risk estimation. For evaluation of the similarity of sampling sites with respect to contribution of metals in urban soils, cluster analysis (CA) was applied¹¹. Hierarchical CA was performed using the Ward's method and Euclidean distances as a measure of similarity and the results are showed in a dendrogram. Before applying CA, the normality of all metals was checked using Shapiro-Wilk's normality test ($p < 0.05$). In this study prior to CA all the data were log-transformed to reduce the influence of high values.

Input parameters (toxicity values) for estimation have been taken USEPAs exposure parameters^{12,13}. Children could be exposed to contaminants from soil *via* three different pathways oral intake ($I_{\text{ingestion}}$), inhalation intake ($I_{\text{inhalation}}$) and through skin exposure (I_{dermal})¹³. Based on this fact noncancer risk assessment in this study was estimated. For intake estimation *via* each exposure pathways the following equitations were used.

$$\text{Intake}_{\text{ingestion}} = \frac{C \times \text{IngR} \times \text{EF} \times \text{ED}}{\text{BW} \times \text{AT}} \times 10^{-6} \quad (1)$$

where, C – concentration of a contaminant in soil (mg/kg), IngR – ingestion rate of soil (mg/day) = 200^{14,15}, EF – exposure frequency (days/year) = 360¹³, ED – exposure duration (years) = 6¹⁶, BW – average body weight (kg) = 20.3¹⁷, AT – averaging time (days) = ED*365¹³

$$\text{Intake}_{\text{inhalation}} = \frac{C \times \text{InhR} \times \text{EF} \times \text{ED}}{\text{PEF} \times \text{BW} \times \text{AT}} \quad (2)$$

where, InhR – inhalation rate (m³/day) = 7.6¹⁸, PEF – particle emission factor = 1.36 × 10⁹ m³/kg¹⁶.

$$\text{Intake}_{\text{dermal}} = \frac{C \times \text{SA} \times \text{SAF} \times \text{ABS} \times \text{EF} \times \text{ED}}{\text{BW} \times \text{AT}} \times 10^{-6} \quad (3)$$

where, SA – surface area of the skin that contacts the soil (cm²) = 2,800¹⁶, SAF – skin adherence factor for soil (mg/cm²) = 0.2¹⁶, ABS – dermal absorption factor (chemical specific) = 0.001 (for all metals)¹⁸⁻²⁰.

In this study, the body weight of 20.3 kg was taken from World Health Organization – WHO reference value¹⁷. After the three exposure pathways intake_{ingestion}, intake_{inhalation} and intake_{dermal} were calculated, hazard quotient (HQ) and HI based on non-cancer toxic risk can be calculated as follows¹³:

$$\text{HQ} = \frac{\text{Intake}}{\text{RfD}} \quad (4)$$

$$\text{HI}_{\text{exP}} = \sum \text{HQ}_{\text{exP}} \quad (5)$$

where, exP are different exposure pathways, respectively.

Reference dose (RfD) (mg/kg/day) is an estimate value of the daily exposure, maximum permissible risk, to the human population, including sensitive subgroups (children) during a lifetime. The values of RfD are showed in Table 1¹⁹.

Table 1

Values for reference doses (RfDs)			
Metals	RfD _{ingestion}	RfD _{dermal}	RfD _{inhalation}
Cu	4E-02	1.2E-02	4E-0
Pb	3.5E-03	5.25E-04	3.5E-03
Cd	1E-03	1E-05	1E-03
Zn	3E-01	6E-02	3E-01

In this study it was assumed that after inhalation, all toxicants bonded to particular matter will have similar health effect as if they are ingested. It was assumed that absorption factor for inhalation and ingestion is 100 and this value was used in this study^{12,19-21}.

Each HQ for different pathways could be calculated and summed to generate HI (Eq.5). If the value of HI < 1, there is no significant risk of noncancerogenic effect. But if the HI > 1, there is probability of occurrence of noncancerogenic effect and it will be increased if HI increases²².

Results

Based on metal concentrations after cluster analysis, urban soils, collected from 31 locations, were classified into two groups and results are presented in Figure 1. The group I con-




Fig. 1 – Dendrogram showing clustering of monitoring sites. (for explanation see under Table 2).

sisted of samples Vr1, Pr2, Pr1, Vr4, Vr5, Vr2, Vr3, Vr9, Vr10, Pr5, Vr8, Pr4, Vr7, Vr6, and Pr3 from locations that were near recreational or residential places. The group II consisted of samples Vr25, Vr26, Vr23, Vr24, Vr21, Vr22, Vr19, Vr20, Vr17, Vr18, Vr15, Vr16, Vr13, Vr14, Vr11, and Vr12 and these sites were near traffic roads and some small building materials facilities such as Vr11 location. Descriptive statistics of the two groups are shown in Table 2. Figure 1 shows the dendrogram of clustering of monitoring sites.

All mean metal concentrations in the group II were higher, except of Cd, compared to the group I. The mean concentration of Pb in the group II was 30% higher than in the group I, while the mean Zn concentration in the group II was 50% higher than in the group I. There is no significant difference between mean concentrations of Cd and Cu in these two groups.

Correlation analysis

Pearson's correlation analysis was applied for each group to analyze the relationships of metal concentrations, and the

results are showed in Table 3. Pb, Cd, Cu and Zn were among significantly positively correlated with each other in the group II. Cu and Zn showed very strong positive correlation (0.85) indicating that also natural source together with traffic and industry contribute to contamination. In the group I, there were no statistically significant correlations among metals, and this might be due to natural content and lower pollution of these sites. Concentrations of Cd and Zn were negatively correlated, but not statistically significant, probably indicating different sources of pollution by these two metals.

The obtained results of noncancerogenic children health risk, based on metal concentrations in urban soils and exposure by three different pathways (ingestion, inhalation and dermal) are shown in Table 4. The results for HI for Pb at all investigated locations are showed in Figure 2. In soil sample at the location Vr11 HI for Pb was 0.8 and it is very close to the upper limit of the safe level¹. HI for Pb (0.68) at the location Vr22, in children was also lower than the upper limit of the safe level¹. HI for Pb at the 29 investigated locations is lower than 0.4. A high Pb concentration in urban soil

Table 2
Metal concentrations of urban soils (mg/kg) in two group sites in Podgorica and different regulations for metal concentrations in urban soil

Groups	Cu		Pb		Cd		Zn	
	(r ± SD)	(min-max)	(r ± SD)	(min-max)	(r ± SD)	(min-max)	(r ± SD)	(min-max)
Group I	51.82±20.0	9.95–99.3	75.99±17.75	2365–10923	2.85 ± 0.60	0.89–3.55	846±5036	15.00–236.54
Group II	53.73±21.18	3.57–87.6	98.19±60.25	41.10–28550	2.77 ± 0.51	1.59–3.65	12559±7781	41.59–383.96
National regulations ²³	100		50		3		300	
Residential/recreational intervention limits	120		100		2		150	

Group I – urban soil samples from locations near recreational and residential places;

Group II – urban soil samples form sites near traffic roads

Table 3
Correlation among concentrations of investigated metals in two group sites of Podgorica

Sites	Cu	Pb	Cd	Zn
Group 2*				
Cu	1	0.58*	0.79 [†]	0.85 [†]
Pb	0.58*	1	0.73 [†]	0.39
Cd	0.79 [†]	0.73 [†]	1	0.69 [†]
Zn	0.85 [†]	0.39	0.69*	1
Group 1*				
Cu	1	0.07	0.44	0.47
Pb	0.07	1	0.10	0.18
Cd	0.44	0.10	1	-0.10
Zn	0.47	0.18	-0.10	1

*for explanation see under Table 2

Table 4
Non-carcinogenic (three exposure pathways) risk for children (95% UCL)

Metal	HQ _{ingestion}			HQ _{inhalation}			HQ _{dermal}			HI _{ex}		
	min	max	mean	min	max	mean	min	max	mean	min	max	mean
Pb	0.07	0.8	0.24	1.8E-06	2.3E-05	7E-06	1.2E-03	1.5E-04	4.5E-03	0.071	0.8	0.25
Cd	8.6E-03	3.5E-02	2.7E-02	2.4E-07	1.0E-06	7.6E-07	2.4E-03	1.0E-02	7.6E-03	0.011	0.05	0.035
Cu	9E-04	2.4E-02	1.3E-02	2.5E-08	7.0E-07	3.5E-07	8.1E-06	2.3E-04	1.2E-04	0.001	0.024	0.015
Zn	6.4E-04	1.2E-02	3.4E-03	1.4E-08	3.5E-07			9.5E-08	7.0E-06		1.7E-04	5.0E-05
	0.0006	0.024			0.004							

HQ – hazard quotient; HI – hazard index; UCL – upper confidence limit.

at the location Vr11 is the result of the vicinity of building materials facility and intensive traffic. Noncancer lead risk (HI) for children at the examined locations of Podgorica is shown in Figure 2.

The major path of children exposure to urban soil that adverse human health by Pb, Cd, Cu and Zn is ingestion, followed by dermal exposure. Contribution of inhalation exposure to HI is the smallest. For noncancer risk HI for chil-




Fig. 2 – Noncancer risk for lead (HI) for children at examined locations of Podgorica, Montenegro.

Group I (Vr1, Pr2, Pr1, Vr4, Vr5, Vr2, Vr3, Vr9, Vr10, Pr5, Vr8, Pr4, Vr7, Vr6, Pr3) – samples from locations that were near recreational and residential places;

Group II (Vr25, Vr26, Vr23, Vr24, Vr21, Vr22, Vr19, Vr20, Vr17, Vr18, Vr15, Vr16, Vr13, Vr14, Vr11, Vr12) – samples from locations that were near traffic rods.

Discussion

Metals concentrations were compared with maximum allowed concentrations (MAC) values, recommended by the National Regulation²³ and the Italian intervention criteria for soils (the residential/ recreational intervention limits fixed by the Italian Environmental Law DM 471/99)²⁴. According to the National Regulation, the mean concentration of Pb in both groups was higher than the prescribed value. The mean concentrations of Cd exceeded the Italian residential/recreational intervention limits. The maximum concentrations of Pb and Cd in urban soils, in both groups exceeded the National Regulation limits and Italian residential/recreational intervention limits. In the group II the maximum Zn values were above the National Regulation limits and residential/recreational intervention limits, while in the group I they were above residential/recreational intervention limits. Higher concentration of all investigated metals in the group II could be explained to its proximity to traffic roads and some industrial locations. Podgorica has been under high urbanization in the past few decades. In the study areas, there were no specific pollution sources of toxic metals, because of that, the toxic metal contamination of the soils was derived from continuous urbanization and development, which can influence human health in the contaminated area. It is important to emphasize that the main road that connects south and east part of Montenegro, goes through the city center, with very intensive and heavy traffic. It is common practice to compare the mean concentration of toxic metals in some urban soils from different urban cities¹. It can be concluded that the existing level of Cd and Cu soil contamination in Podgorica is significantly higher than comparable levels in some other cities over the world. Zinc concentrations vary from city to city, while Pb content is the lowest except in the city of Madrid, Spain.

dren decreases in order Pb > Cd > Cu > Zn. So, Pb exposure to the urban soil in Podgorica may also pose health threat, specially, to young children. The daily ingestion rates of soil by children was calculated to be between 39 and 270 mg/day²⁵. Because of its negative effects on the children's central nervous system, monitoring of Pb content in soil is of great importance²⁶. Many neurological and developmental disorders may be observed in children's population due to the long period of exposure and ingestion of certain amounts of Pb from contaminated soil, such as anemia, kidney damage, colic, muscle weakness and brain damage^{19, 21, 27}. Ingestion of small quantity of Pb from dust may be harmful for blood, development, behavior and intellectual functioning, as well^{27, 28}.

Because of such threats to children's health it is necessary to take action to decontaminate locations where HI is below the safe level, but high enough to adverse children's health during a long-time exposure. The potential health risk from Cd and Cu is low while potential health risk from Zn is the least for children's population.

Conclusion

Based on the content of toxic metals, Pb, Cd, Cu and Zn, in urban soils, playgrounds and parks, in Podgorica, Montenegro and after cluster analysis all metals showed relatively higher concentrations at sites that were close to industrial places and highways clearly indicating the influence of rapid urbanization and industrialization in the last few decades. The mean concentration of Pb and maximum concentrations of Pb, Cd and Zn were higher than the prescribed value in the National Regulation. The highest risk is associated with soil particle ingestion and the noncarcinogenic health risk for children was very high at two locations. Because of such threats to chil-

dren's health it is necessary to take action to decontaminate these sites and prevent and protect children's health. This integrated approach based on statistical and non-carcinogenic probabilistic risk analysis may help in the decision making process in every growing urban and industrial region.

Acknowledgement

The authors wish to thank the Ministry of Science and Ministry of Health of Montenegro for financial support through the Project No. 03-401.

R E F E R E N C E S

1. Marjanović MD, Vukčević MM, Antonović DG, Dimitrijević SI, Jovanović DM, Mataušić M, et al. Heavy metals concentration in soils from parks and green areas in Belgrade. *J Serb Chem Soc* 2009; 74(6): 697–706.
2. Abrahams PW. Soils: their implications to human health. *Sci Total Environ* 2002; 291(1–3): 1–32.
3. Daydova S. Heavy metals as toxicants in big cities. *Microchem J* 2005; 79(1–2): 133–316.
4. Poggio L, Vrsčaj B, Schulín R, Hepperle E, Ajmone Marsan F. Metals pollution and human bioaccessibility of topsoils in Grugliasco (Italy). *Environ Pollut* 2009; 157(2): 680–9.
5. Imperato M, Adamo P, Naimo D, Arienzo M, Stanzione D, Violante P. Spatial distribution of heavy metals in urban soils of Naples city (Italy). *Environ Pollut* 2003; 124(2): 247–56.
6. Ljung K, Selinus O, Otabbong E. Metals in soils of children's urban environments in the small northern European city of Uppsala. *Sci Tot Environ* 2006; 366(2–3): 749–59.
7. Acosta JA, Cano AF, Arocena JM, Debela F, Martínez-Martínez S. Distribution of metals in soil particle size fractions and its implication to risk assessment of playground in Murcia City (Spain). *Geoderma* 2009; 149(1–2): 101–9.
8. Popoola OE, Bamgbose O, Okonkwo OJ, Aronwo TA, Odukoya AT, Popoola AO. Heavy Metals Content in Playground Topsoil of Some Public Primary Schools in Metropolitan Lagos, Nigeria. *Res J Environ Earth Sci* 2012; 4(4): 434–9.
9. Tong ST, Lam KC. Home sweet home? A case study of household dust contamination in Hong Kong. *Sci Tot Environ* 2000; 256(2–3): 115–23.
10. Nriagu JO. A silent epidemic of environmental metal poisoning. *Environ Pollut* 1988; 50(1–2): 139–61.
11. Chabukdhara M, Nema AK. Heavy metals assessment in urban soil around industrial clusters in Ghaziabad, India: probabilistic health risk approach. *Ecotoxicol Environ Saf* 2013; 87: 57–64.
12. US EPA (United States Environmental Protection Agency). Risk Assessment Guidance for Superfund. Human Health Evaluation Manual. EPA/540/1-89/002, vol. I. Office of Solid Waste and Emergency Response. 1989. Available from: <http://www.epa.gov/superfund/programs/risk/ragsa/index.htm>
13. US DoE. The Risk Assessment Information System (RAIS). Argonne, IL: U.S. Department of Energy's Oak Ridge Operations Office (ORO); 2011.
14. Calabrese EJ, Kosteckí PT, Gilbert CE. How much dirt do children eat?, An emerging environmental health question. *Comments Toxicol* 1987; 1: 229–41.
15. US EPA (United States Environmental Protection Agency). Exposure Factors Handbook. EPA/600/P-95/002F. Washington, DC: Environmental Protection Agency, Office of Research and Development; 1997.
16. US EPA (United States Environmental Protection Agency). Supplemental Guidance for Developing Soil Screening Levels for Superfund Sites. OSWER 9355.4-24.2001. Office of Solid Waste and Emergency Response. 2001. Available from: <http://www.epa.gov/superfund/resources/soil/ssgmarch01.pdf>
17. World Health Organization. Growth reference data for 5-19 years. Geneva: World Health Organization; 200. Available from: <http://www.who.int/growthref/en>
18. Zheng N, Liu J, Wang Q, Liang Z. Health risk assessment of heavy metal exposure to street dust in the zinc smelting district, northeast of China. *Sci Total Environ* 2010; 408(4): 726–33.
19. de Miguel E, Iribarren I, Chacón E, Ordoñez A, Charlesworth S. Risk-based evaluation of the exposure of children to trace elements in playgrounds in Madrid (Spain). *Chemosphere* 2007; 66(3): 505–13.
20. Shi G, Chen Z, Bi C, Wang L, Ten J, Li Y, et al. A comparative study of health risk of potentially toxic metals in urban and suburban road dust in the most populated city of China. *Atmos Environ* 2011; 45(3): 764–71.
21. Ferreira-Baptista L, de Miguel E. Geochemistry and risk assessment of street dust in Luanda, Angola: a tropical urban environment. *Atmos Environ* 2005; 39(25): 4501–12.
22. Lee CS, Li XD, Shi WZ, Cheung SC, Thornton I. Metal contamination in urban, suburban, and country park soils of Hong Kong: a study based on GIS and multivariate statistics. *Sci Total Environ* 2006; 356(1–3): 45–61.
23. Rulebook on permitted amounts of hazardous and noxious substances in soil and methods for their testing. Official Gazette of Montenegro No 18; 1997.
24. Italian Ministry of Environment. Technical Regulation D.M. No. 471/1999 on containment, remediation and environmental restoration of contaminated soils. Annex of the Italian Official Gazette No. 293, 15.12.1999. (Italian)
25. Ljung K, Oomen A, Duits M, Selinus O, Berglund M. Bioaccessibility of metals in urban playground soils. *J Environ Sci Heal* 2007; 42(9): 1241–50.
26. Cicchella D, de Vivo B, Lima A, Albanese S, McGill RA, Parrish RR. Heavy metal pollution and Pb isotopes in urban soils of Napoli, Italy. *Geochem Expl Environ Anal* 2008; 8(1): 103–12.
27. Osman K. Health effects of environmental lead exposure in children [dissertation]. Stockholm: Karolinska Institute, Institute of Environmental Medicine; 1998. (Swedish)
28. ATSDR. Public Health Statements - Toxicological Profiles (Al, As, Cd, Co, Cr, Cu, Hg, Mn, Ni, Pb, Zn). 2002. [online] [updated 2006 March 14] Available from: <http://www.atsdr.cdc.gov/phshome.html>

Received on July 16, 2013.

Revised on June 9, 2014.

Accepted on September 11, 2014.

Online First July, 2015.