



The Balkan Botanical Congress is an international meeting that has been held nearly every three years, since 1997. It brings together botanists from around the world who perform research on plants in the widest sense, as well as scientists who are engaged in the plant sciences and their applications. We were honored to host such an extraordinary scientific event this year in Serbia.

The 7th Balkan Botanical Congress – 7BBC 2018 took place in Novi Sad from September 10th to 14th 2018. The Congress was organized by the University of Novi Sad, Faculty of Sciences, Department of Biology and Ecology and the “Andreas Wolny” Botanical Society, along with the great help of 7 co-organizers and more than 30 supporters and sponsors. It truly was not possible to happen without exceptional help of our co-organizer - the Institute for Nature Conservation of Vojvodina Province who made this congress not only possible, but totally awesome.

7BBC 2018 placed a special emphasis on plants of the Balkan Peninsula and covered various research fields. The Congress was organized into ten sessions: Plant Anatomy and Physiology, Plant Taxonomy and Systematics, Plant Molecular Biology and Genetics, Floristics, Vegetation and Phytogeography, Conservation Botany and Plant Invasions, Phytochemistry and Plant Resources, Agronomy and Forestry, Botanical Collections and History, Ethnobotany and Cryptogam Biology. These topics were elaborated through five plenary lectures given by eminent scientists, as well as in the form of introductory lectures, oral and poster presentations. With an overall number of 387 abstracts presented on the very latest of botanical science, we shared knowledge, expertise and novel ideas. We welcomed nearly 400 scientists to Novi Sad, and we believe that we succeeded in our joint endeavor to make new networks and new connections among botanists. We hope that we contributed to advancements in the wide and beautiful field of botany, ranging from fundamental botanical research to applied botany.

It is our great pleasure to publish this Abstract Book in Botanica Serbica, in the same year that this international journal, a renamed continuation of the Bulletin of the Institute of Botany and Botanical Garden Belgrade, celebrates its 90 year jubilee. On behalf of the Scientific and Organizing committee of 7BBC 2018 we would like to express our gratitude to all contributors, colleagues and sponsors for taking part in the 7th Balkan Botanical Congress, as well as for their efforts and contributions to its successful realization.

Goran Anačkov and Lana Zorić,
Co-presidents of the Scientific Committee of the 7 BBC
and guest editors of Botanica Serbica 42 (supplement 1).

Organizers:

University of Novi Sad, Faculty of Sciences, Department of Biology and Ecology, Novi Sad
Botanical Society „Andreas Wolny“, Novi Sad

Co-organizers:

Institute for Nature Conservation of Vojvodina Province, Novi Sad
Institute for Nature Conservation of Serbia, Belgrade
University of Belgrade, Faculty of Biology, Belgrade
University of Belgrade, Faculty of Forestry, Belgrade
University of Belgrade, Institute for Biological Research “Siniša Stanković“, Belgrade
University of Novi Sad, Faculty of Medicine, Center for Medical-Pharmaceutical Research and Quality Control, Novi Sad
Natural History Museum in Belgrade, Belgrade

Support:

Republic of Serbia, Ministry of Education, Science and Technological Development
Republic of Serbia, Ministry of Environmental Protection
Republic of Serbia, Autonomous Province of the Vojvodina, Provincial Secretary for Higher Education and Scientific Research Activity
Republic of Serbia, Autonomous Province of the Vojvodina, Provincial Secretary for Urbanization and Environmental Protection
City of Novi Sad
PWMC “Vode Vojvodine“, Novi Sad
PC “Vojvodinašume“, Petrovaradin
PCC “Gradsko zelenilo“, Novi Sad
PCC “Lisje“, Novi Sad
Matica srpska, Novi Sad
Institute of Field and Vegetable Crops, Novi Sad
University of Novi Sad, Institute of Lowland Forestry and Environment, Novi Sad
University of Novi Sad, Institute of Food Technology in Novi Sad, Novi Sad
University of East Sarajevo, Faculty of Technology, Zvornik
Journal “Plant Systematics and Evolution“
World Wild Fund For Nature, Belgrade
IUCN ECARO, Belgrade
Vojvodina Environmental Movement, Novi Sad
Biology and Ecology Students’ Scientific Research Society “Josif Pančić“, Novi Sad
National Park “Fruška gora”
Nature Park “Rusanda”
SNR “Deliblato Sand”
SNR “Obedska bara”
SNR “Okanj bara”
SNR “Slano Kopovo”
SNR “Titelski breg”
SNR “Zasavica”
Hungarian Natural History Museum, Budapest
Tourism Organization of Vojvodina
Tourist Organization of the City of Novi Sad, Novi Sad
PanaComp, Wonderland Travel, Novi Sad

Sponsors:

- Coca-Cola HBC, Belgrade
- Naftachem, Sremski Karlovci
- BioSPIN ltd, Novi Sad
- Mikronik ltd, Belgrade

- Nikon
- Pivnica “Gusan“, Novi Sad
- Intercaffe ltd, Belgrade

Honorable Commitee

Dr Ana Petrova, Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Bulgaria
Dr Kit Tan, Department of Biology, Faculty of Science, University of Copenhagen, Denmark
Dr Arne Strid, Department of Biology, Faculty of Science, Lund University, Sweden
Dr Werner Greuter, Herbarium Mediterraneum, University of Palermo, Italy & Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie University of Berlin, Germany
Dr Branislava Butorac, Institute for Nature Conservarion, Serbia
Dr Branka Stevanović, Faculty of Biology, University of Belgrade, Serbia
Dr Dušan Nikolić, Rector of University of Novi Sad, Serbia
Dr Jelena Blaženčić, Faculty of Biology, University of Belgrade, Serbia
Dr Milica Pavkov Hrvojević, Dean of Faculty of Sciences, University of Novi Sad, Serbia
Miloš Vučević, The Mayor of Novi Sad, Serbia
Dr Pal Boža, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
Dr Rudolf Kastori, Secretary General of the Department of Natural Sciences, Matica srpska, Serbia
Dr Vladimir Stevanović, Faculty of Biology, University of Belgrade & Serbian Academy of Sciences and Art, Serbia
Vladimir Galić, Provincial Secretary for Urban Planning and Environmental Protection, Serbia
Dr Zoran Milošević, Provincial Secretary for Higher Education and Scientific Research, Serbia
Dr Karol Marhold, Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Charles University, Prague, and Secretary-General of International Association for Plant Taxonomy, Slovak Republic & Czech Republic
Dr Tod Stuessy, Museum of Biological Diversity, The Ohio State University, United States of America

Scientific Committee**Presidents:**

Dr Goran Anačkov, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
Dr Lana Zorić, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia

Members:

Dr Alfred Mullaj, Faculty of Natural Sciences, University of Tirana, Albania
Dr Lulëzim Shuka, Department of Biology, Faculty of Natural Sciences, University of Tirana, Albania
Dr Božo Frajman, Institute of Botany, University of Innsbruck, Austria
Dr Peter Schönswetter, Institute of Botany, University of Innsbruck, Austria
Dr Faruk Bogunić, Faculty of Forestry, University of Sarajevo, Bosnia and Herzegovina
Dr Senka Barudanovic, Faculty of Science, Bosnia and Herzegovina
Dr Siniša Škondrić, Department of Biology, Faculty of Sciences, University of Banja Luka, Bosnia and Herzegovina
Dr Rosen Tsonev, Faculty of Biology, Sofia University “St. Kliment Ohridski“, Bulgaria
Dr Vladimir Vladimirov, Department of Plant and Fungal Diversity and Resources, Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Bulgaria
Dr Antun Alegro, Department of Biology, Faculty of Science, University of Zagreb, Croatia
Dr Boštjan Surina, Natural History Museum Rijeka, Croatia
Dr Sandra Bogdanović, Faculty of Agriculture, University of Zagreb, Croatia
Dr Sonja Šiljak Jakovljević, Ecologie Systématique Evolution, CNRS, AgroParisTech, Univ. Paris-Sud, Université Paris-Saclay, France
Dr Dimitris Tzanoudakis, Division of Plant Biology, Department of Biology, University of Patras, Greece
Dr Panayotis Dimopoulos, Institute of Botany, Division of Plant Biology, Department of Biology, University of Patras, Greece
Dr Theophanis Constantinidis, Department of Ecology and Systematics, Faculty of Biology, National and Kapodistrian University of Athens, Greece
Dr Király Gergely, Institute of Silviculture and Forest Protection, University of Sopron, Hungary
Dr Zoltán Barina, Department of Botany, Hungarian Natural History Museum, Hungary
Dr Vlado Matevski, Institute of Biology, Faculty of Natural Sciences and Mathematics, Ss. Cyril and Methodius University and Macedonian Academy of Sciences and Arts, Macedonia
Dr Danka Caković, Faculty of Natural Sciences and Mathematics, University of Montenegro, Montenegro

Dr Danijela Stešević, Faculty of Natural Sciences and Mathematics, University of Montenegro, Montenegro
 Dr Vesna Mačić, Institute of Marine Biology, University of Montenegro, Montenegro
 Dr Łuczaj Łukasz, Department of Botany, Institute of Applied Biotechnology and Basic Sciences, University of Rzeszów, Poland
 Dr László Bartha, Institute for Interdisciplinary Research in Bio-Nano Sciences Romania
 Dr Biljana Božin, Department of Pharmacy, Faculty of Medicine, University of Novi Sad, Serbia
 Dr Bojan Konstantinović, Department of Environmental and Plant Protection Faculty of Agriculture, University of Novi Sad, Serbia
 Dr Bojan Zlatković, Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš, Serbia
 Dr Branislava Lakušić, Faculty of Pharmacy, University of Belgrade, Serbia
 Dr Dmitar Lakušić, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Dragana Miladinović, Institute of Field and Vegetable Crops, Serbia
 Dr Dragana Rančić, Faculty of Agriculture, University of Belgrade, Serbia
 Dr Dragana Vukov, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Gordana Tomović, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Ivana Maksimović, Faculty of Agriculture, University of Novi Sad, Serbia
 Dr Jadranka Luković, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Maja Karaman, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Marjan Niketić, Natural History Museum, Serbia
 Dr Marko Sabovljević, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Mihajla Đan, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Milan Stanković, Department of Biology and Ecology, Faculty of Sciences, University of Kragujevac, Serbia
 Dr Milan Veljić, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Mirjana Šijačić Nikolin, Faculty of Forestry, University of Belgrade, Serbia
 Dr Miroslava Mitrović, Institute for Biological Research “Siniša Stanković”, University of Belgrade, Serbia
 Dr Nataša Nikolić, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Neda Mimica Dukić, Department of Chemistry, Biochemistry and Environmental Protection, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Pavle Pavlović, Institute for Biological Research “Siniša Stanković”, University of Belgrade, Serbia
 Dr Peđa Janačković, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Petar Marin, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Saša Orlović, Institute of Lowland Forestry and Environment, University of Novi Sad, Serbia
 Dr Slobodan Jovanovic, Institute of Botany and Botanical Garden “Jevremovac”, Faculty of Biology, University of Belgrade, Serbia
 Dr Slobodanka Pajević, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Snežana Radulović, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Serbia
 Dr Srđan Stojnić, Institute of Lowland Forestry and Environment and Faculty of Agriculture, University of Novi Sad, Serbia
 Dr Vladimir Ranđelović, Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš, Serbia
 Dr Andraž Čarni, “Jovan Hadži” Institute of Biology, Slovenia
 Dr Nejc Jogan, Biotechnical Faculty, University of Ljubljana, Slovenia
 Dr Neriman Özhatay, Department Of Pharmaceutical Botany, Faculty of Pharmacy, Istanbul University, Turkey

Organizing Committee

Presidents:

Dr Ružica Igić, President of Botanical Society “Andreas Wolny”, Novi Sad
 Dr Biljana Panjković, Head of Institute for Nature Conservation of the AP Vojvodina, Novi Sad

Secretaries:

Bojana Bokić, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Milica Rat, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad

Members:

Dr Biljana Božin, Department of Pharmacy, Faculty of Medicine, University of Novi Sad
 Dr Dragana Vukov, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Dušanka Cvijanović, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Goran Anačkov, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Jadranka Luković, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Lana Zorić, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Ljiljana Nikolić, Faculty of Agriculture, University of Novi Sad

Dr Milan Borišev, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Milan Župunski, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dr Nebojša Kladar, Department of Pharmacy, Faculty of Medicine, University of Novi Sad
 Dr Slobodanka Pajević, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Ana Vestek, Botanical Society “Andreas Wolny” Novi Sad
 Boris Radak, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Danijela Arsenov, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Dragan Obradov, Botanical Society “Andreas Wolny” Novi Sad
 Dunja Karanović, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Đurđica Simin, Botanical Society “Andreas Wolny” Novi Sad
 Goran Tmušić, Botanical Society “Andreas Wolny” Novi Sad
 Jelena Jocković, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Jelena Knežević, Botanical Society “Andreas Wolny” Novi Sad
 Marija Kovački, Botanical Society “Andreas Wolny” Novi Sad
 Marko Ručando, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Miloš Ilić, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Mirjana Čuk, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad
 Ranko Perić, Institute for Nature Conservation of the AP Vojvodina
 Sara Pavkov, Institute for Nature Conservation of the AP Vojvodina
 Slobodan Bojčić, Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad

Sessions:

The 7th Balkan Botanical Congress consists of plenary lectures, introductory lectures of each session, as well as oral and poster presentations on the following topics:

Sessions 1. Plant Anatomy and Physiology

Sessions 2. Plant Taxonomy and Systematics

Sessions 3. Plant Molecular Biology and Genetics

Sessions 4. Floristics, Vegetation and Phytogeography

Sessions 5. Conservation Botany and Plant Invasion

Sessions 6. Phytochemistry and Plant Resources

Sessions 7. Agronomy and Forestry

Sessions 8. Botanical Collections and History

Sessions 9. Ethnobotany

Sessions 10. Cryptogam Biology

in cortex, some are just below the epidermis and many, that are in forming phase above the phloem. The older stem (perennial stem) has many secretory canals in the parenchyma of cortex and in the xylem of secondary wood. The essential oils were obtained by hydrodistillation and qualitative and quantitative analysis was performed by GC-FID and GC-MS. Small amounts of essential oils were extracted from the needles (0.06-0.41%), stems (0.31-0.54%) and strobilus (0.16 % and 0.19 %) of *L. decidua*. The monoterpene (28.70-50.40%) and sesquiterpene compounds (41.80-67.20%) dominated in the essential oil of needles, monoterpenes (80.50-91.70%) in the oil of stems, diterpenes (46.50% and 40.50%) and monoterpenes (24.20% and 22.20%) in the oil of strobilus. Main compounds in the essential oil of needles represented germacrene D (15.2-46.9%), δ -3-carene (4.6-24.6%) and α -pinene (6.3-14.1%), in the oil of stems α -pinene (12.8-21.8%), δ -3-carene (8.6-14.3%) and sabinene (0.5-10.3%), while in the oil of strobilus diterpene of abietane type (4.1-10.2%) and nor-abietatriene (3.9-9.2%) were the most abundant. The larch needles and stem essential oils from cultivated and natural habitats were quite similar in qualitative composition. However, the essential oil of needles from natural habitat contained higher content of germacrene D and α -pinene and the oil of stem α -pinene and trans-verbenaol. The oil of larch strobilus from natural habitat was characterized by manool and 3- α -acetoxy-manool, diterpenes that were not present in the oil of cultivated sample.

KEYWORDS: *Larix decidua*, anatomy, needles, stem, strobilus, essential oil

Poster presentation 10 06 53

PIMPINELLA TRAGIUM VILL. (APIACEAE) – ANATOMY AND ESSENTIAL OIL COMPOSITION

Violeta Slavkovska^{1*}, Bojan Zlatković², Nada Kovačević³ & Milica Drobac³

¹Department of Botany, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, 11221 Belgrade, Serbia, ²Department of Biology and Ecology, Faculty of Sciences and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia, ³Department of Pharmacognosy, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, 11221 Belgrade, Serbia

*Corresponding author: violetas@pharmacy.bg.ac.rs

Pimpinella tragiium Vill. is a perennial, herbaceous plant inhabiting dry grasslands, limestone rocky areas and screes of wide sense Mediterranean area, east European lowlands and Caucasus. This study explores the anatomical structure, content and composition of the essential oil of the vegetative organs and fruit of this species. The plant material was collected on two localities: the village Izvor (Bosilegrad, Serbia) and Mt. Galičica (FYR Macedonia). The anatomical studies were conducted on permanent slides obtained by standard

method of preparation for viewing under a light microscope. The essential oils obtained by hydrodistillation from roots (collected during flowering and fruiting period), aerial parts with inflorescences, and fruits, were analysed by GC-FID and GC-MS. The anatomical analysis revealed a secondary structure of root and primary structure of stem with closed collateral vascular bundles. Leaves are isobilateral, amphistomatic; petiole is with arched vascular bundles and the fruit (mericarp) is semi-circular and lightly ribbed in cross section. Non-glandular unicellular and bicellular cuneate trichomes are sparsely distributed on the stem, leaves and petiole, but very dense on the fruit. Secretory channels are present in all organs: in the parenchyma of the root and stem cortex, stem pith, in the phloem of root and vascular bundles of stem and petiole, by the leaf vascular bundles and in the fruit pericarp. Essential oil yields from different parts of *P. tragiium* ranged from 0.2-1.1% (v/w). The main compounds in the oils from roots (in both stages, from both localities), as well as from aerial parts and fruits from Mt. Galičica are C-12 norsesquiterpenes (trinorsesquiterpenes): pregeijerene (29.0-56.2%) and gejerene (14.1-22.9%), whereas those from aerial parts and fruits from village Izvor are β -bisabolene (19.1-57.2%) and a phenylpropanoid epoxy-pseudoisoeugenyl-2-methoxybutyrate (17.4-22.2%). Trinorsesquiterpenes, which were also found to be characteristic constituents in some other *Pimpinella* species oils, are the most dominant constituents in all investigated oils, except in the oil of fruit from village Izvor. In addition, phenylpropanoids of pseudoisoeugenol type are present in all oils confirming them as chemical markers of *Pimpinella* species analysed so far.

KEYWORDS: *Pimpinella tragiium*, anatomy, secretory canals, essential oil, C-12 norsesquiterpenes, phenylpropanoids

Poster presentation 11 06 29

CHEMOSYSTEMATIC EVALUATION OF THE COMPOSITION OF LEAF AND FLOWER ESSENTIAL OILS OF EIGHT HERACLEUM L. TAXA FROM SOUTHEASTERN EUROPE

Ljuboš Ušjak^{1*}, Milica Drobac¹, Marjan Niketić² & Silvana Petrović¹

¹Department of Pharmacognosy, University of Belgrade - Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia, ²Natural History Museum, Njegoševa 51, 11000 Belgrade, Serbia

*Corresponding author: ljubos@pharmacy.bg.ac.rs

The genus *Heracleum* L. (Apiaceae) is represented by a dozen of native sub(species) in Southeastern Europe. In this study, the composition of the essential oils obtained from the leaves and the flowers of eight taxa of this genus, collected in Serbia, Montenegro, Macedonia and Slovenia, was statistically analyzed to evaluate its chemosystematic significance. Investigated taxa included *H. orphanidis* Boiss. and the representatives of *H. sphondylium* group: *H. sphondylium* L., *H. sibiricum* L., *H. montanum* Schleich. ex Gaudin, *H. ternatum* Velen., *H. pyrenaicum* subsp. *pollinianum* (Bertol.) F. Pedrotti & Pignatti

ti, *H. pyrenaicum* subsp. *orsinii* (Guss.) F. Pedrotti & Pignatti and *H. verticillatum* Pančić. Essential oils were hydrodistilled using Clevenger-type apparatus and analyzed by GC-FID and GC-MS. Chemosystematic significance of their components was evaluated using multivariate statistics: principal component analysis (PCA), non-metric multidimensional scaling (nMDS) and unweighted pair-group arithmetic averages clustering (UPGMA). The analyses included our previously published data on the oils of eight samples of the leaves and three of the flowers, as well as additionally analyzed oils of eight samples of the leaves and five of the flowers. Leaf and flower oils of investigated members of *H. sphondylium* group were dominated by various sesquiterpenes [(*E*)-caryophyllene, (*E*)-nerolidol, (*E*)- β -farnesene, α -trans-bergamotene, germacrene D, β -bisabolene and/or β -sesquiphellandrene], phenylpropanoids [apiol, methyl eugenol, elemicin and/or (*Z*)-isoelemicin], and/or monoterpene limonene. On the other hand, leaf and flower oils of *H. orphanidis* were rich in aliphatic esters, mostly octyl acetate. Separate statistical analyses of the compositions of the leaf oils and the flower oils demonstrated segregation of *H. orphanidis* from investigated representatives of *H. sphondylium* group, and grouping of the subspecies of *H. pyrenaicum* within this group. Morphologically related species *H. sibiricum* and *H. ternatum* were closely located in PCA and nMDS, and in UPGMA even shared the same cluster. PCA showed that some of both aforementioned dominant constituents and those present in lower amounts influenced the separation of investigated taxa. It can be concluded that applied multivariate statistical methods demonstrated the grouping of investigated *Heracleum* taxa according to their current systematics, and justify further similar study on the essential oils of more species of this genus.

KEYWORDS: *Heracleum* taxa, leaf and flower essential oils, GC-FID and GC-MS, PCA, nMDS, UPGMA

Poster presentation 12 06 30

DPPH RADICAL SCAVENGING POTENTIAL OF THE ROOT ESSENTIAL OILS OF FIVE HERACLEUM L. TAXA

Ljuboš Ušjak^{1*}, Milica Drobac¹, Marjan Niketić² & Silvana Petrović¹

¹Department of Pharmacognosy, University of Belgrade - Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia, ²Natural History Museum, Njegoševa 51, 11000 Belgrade, Serbia

*Corresponding author: ljubos@pharmacy.bg.ac.rs

In the Balkan Peninsula, *Heracleum* L. taxa (Apiaceae) were traditionally used for the treatment of various digestive and respiratory diseases, epilepsy, hypertension and sexual weakness. The purpose of this work was to investigate 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging ability of the root essential oils of some of the Balkan *Heracleum* taxa, i.e. *H. sibiricum* L., *H. ternatum* Velen., *H. verticillatum* Pančić,

H. pyrenaicum subsp. *pollinianum* (Bertol.) F. Pedrotti & Pignatti and *H. pyrenaicum* subsp. *orsinii* (Guss.) F. Pedrotti & Pignatti. GC-FID and GC-MS analysis of these oils, obtained by hydrodistillation using Clevenger-type apparatus, revealed the domination of monoterpenes, mostly β -pinene (26.2-47.3%). Additionally, *H. sibiricum* root oil was rich in phenylpropanoids, mainly elemicin (25.6%) and methyl eugenol (22.3%). In colorimetric DPPH assay, the strongest activity was exhibited by *H. sibiricum* oil (SC_{50} =5.19 μ L/mL), followed by *H. pyrenaicum* subsp. *orsinii*, *H. ternatum*, *H. pyrenaicum* subsp. *pollinianum* and *H. verticillatum* oils (SC_{50} =7.85-12.33 μ L/mL). In TLC-DPPH (dot-blot) test, three the most active root oils, i.e. those of *H. sibiricum*, *H. ternatum* and *H. pyrenaicum* subsp. *orsinii* revealed yellow anti-DPPH zones (R_f =0.30-0.42), which were then eluted and analyzed by GC-FID and GC-MS. It was shown that elemicin and methyl eugenol, dominant in *H. sibiricum* oil, were also the most abundant in its anti-DPPH zone (64.5 and 19.5%). β -Pinene and other monoterpene hydrocarbons were not detected in this, and also in the active zones of *H. ternatum* and *H. pyrenaicum* subsp. *orsinii* oils (both contained two closely located anti-DPPH zones, which were eluted together). Anti-DPPH zones of *H. ternatum* oil were dominated by trans-sabinol (21.8%), spathulenol (21.7%) and (*E*)-sesquilandulol (13.1%), and those of *H. pyrenaicum* subsp. *orsinii* oil by (*E*)-sesquilandulol (14.5%) and intermedeol (13.6%). These oxygenated terpenes were detected in *H. ternatum* and *H. pyrenaicum* subsp. *orsinii* oils only in small quantities. Another minor constituent of *H. pyrenaicum* subsp. *orsinii* root oil, (*Z*)-falcarinol, was among the dominant ones in its anti-DPPH zones (11.6%).

KEYWORDS: *Heracleum* taxa, root essential oils, GC-FID and GC-MS, DPPH radical, phenylpropanoids, oxygenated terpenes

Poster presentation 13 06 36

CHEMICAL COMPOSITION OF ESSENTIAL OIL OF ENDEMIC SPECIES ACINOS ORONTIUS FROM BOSNIA AND HERZEGOVINA

Danijela Vidic^{1*}, Amira Čopra-Janičijević¹, Nermina Korać^{1,2} & Milka Maksimović¹

¹Department of Chemistry, Faculty of Science, University of Sarajevo, Sarajevo, Bosnia and Herzegovina, ²Institute of Occupational Health Sarajevo, Laboratory for Toxicology Studies and Sanitary Work Environment Sarajevo, Bosnia and Herzegovina

*Corresponding author: danijela.vidic@gmail.com

The genus *Acinos* belongs to the family Lamiaceae, and represented by ten species native to southern Europe and western Asia. Its name comes from the Greek word *akinos*, the name of a small aromatic plant. *Acinos orontius* (K. Malý) Šilić is a synonym of *Clinopodium alpinum* subsp. *orontium* (K. Malý) Govaerts, and is endemic species in Bosnia and Herzegovina. The aim of this work was to determine content and composi-